

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/259317357>

Trend in the prevalence of obesity and overweight among Iranian children and adolescents: A systematic review and meta-analysis

Article in *Nutrition* · April 2014

DOI: 10.1016/j.nut.2013.08.011 · Source: PubMed

CITATIONS

31

READS

426

4 authors:



[Roya Kelishadi](#)

Isfahan University of Medical Sciences

555 PUBLICATIONS 7,054 CITATIONS

[SEE PROFILE](#)



[AliAkbar Haghdoost](#)

Kerman University of Medical Sciences

375 PUBLICATIONS 2,267 CITATIONS

[SEE PROFILE](#)



[Behnam Sadeghirad](#)

McMaster University

36 PUBLICATIONS 339 CITATIONS

[SEE PROFILE](#)



[Razieh Khajekazemi](#)

Kerman University of Medical Sciences

16 PUBLICATIONS 121 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:




Estimating the prevalence of high-risk behaviors using network scale-up method in university students of Larestan in 2014 [View project](#)



Forecasting Iran Curative Resources Needs for 2025 [View project](#)

All content following this page was uploaded by [Behnam Sadeghirad](#) on 16 November 2014.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.



Review

Trend in the prevalence of obesity and overweight among Iranian children and adolescents: A systematic review and meta-analysis

Roya Kelishadi M.D.^a, Ali-Akbar Haghdoost M.D., Ph.D.^b,
Behnam Sadeghirad PharmD., M.P.H.^c, Razieh Khajehkazemi M.Sc.^{d,*}

^a Child Growth and Development Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

^b Research Center for Health Services Management, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

^c Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran

^d Research Center for Modeling in Health, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

ARTICLE INFO

Article history:

Received 11 May 2013

Accepted 17 August 2013

Keywords:

Obesity

Overweight

Children

Iran

Systematic review

Meta-analysis

ABSTRACT

Objective: Childhood obesity is an important predisposing factor for most non-communicable diseases. The aim of this review was to provide evidence on the prevalence and trends of childhood obesity and overweight in Iran.

Methods: Multiple international and Iranian scientific databases were searched for relevant literatures. Two independent reviewers identified relevant papers in several steps. Separate meta-analyses (using fixed- or random-effect models) were performed to estimate the overall, age, sex, and age–sex specific prevalence of obesity and overweight. Stratified analysis based on Centers for Disease Control and Prevention, International Obesity Task Force, and World Health Organization definition criteria and study year also were performed.

Results: We included 107 studies in the meta-analysis (49 English and 58 Persian). Based on Centers for Disease Control and Prevention definition criteria, the overall prevalence of obesity and overweight remained relatively constant in the 2000s and are estimated to be about 5.1% (95% confidence interval [CI], 4.4–5.8) and 10.8% (95% CI, 10.2–11.4), respectively. The meta-regression analysis showed that the prevalence of obesity and overweight did not vary significantly with respect to sex and age of study participants. Girls had a lower prevalence of obesity and higher prevalence of overweight than boys.

Conclusion: This review, which is the first of its kind in the Middle East and North Africa, suggests that although the trend in the prevalence of childhood obesity in Iranian children is not considerably high, but the escalating trend of excess weight among young children is alarming and should be considered by providers of interventional preventive programs at national and regional levels.

© 2014 Elsevier Inc. All rights reserved.

Introduction

The epidemic of non-communicable diseases (NCD) is a global problem that is expected to rise considerably in low- and middle-income countries [1,2]. Excess weight is one of the most common risk factors for most NCDs. A growing body of evidence exists about the worldwide increase in mean body mass index

(BMI) [3], as well as the prevalence of obesity and overweight [4], with substantial variation in levels and trends in different countries. Excess weight is not limited to adults, and it has become an emerging health problem in the pediatric age group, even in developing countries [5,6].

Of special concern in this regard is the situation in the Middle East and North Africa (MENA), which is facing a double burden of nutritional disorders. The epidemiologic transition, lifestyle changes, and considerably high prevalence of obesity has led the Middle Eastern population to face the greatest global burden of NCDs, notably diabetes [7] and cardiovascular diseases [8]. As a country in this region, Iran has a high prevalence of obesity and

The literature search, data extraction and analysis were done by RK and BS. All authors contributed equally to development of search strategy, interpreting the results, and development of paper. The authors report no conflict of interest.

* Corresponding author. Tel.: +98 341 226 3725; fax: +98 341 226 3725.

E-mail address: r.khajehkazemi@gmail.com (R. Khajehkazemi).

related disorders, such as metabolic syndrome and diabetes in different age groups [9–11]. Such disorders pose high costs to the health care system. For instance, the total national cost of diagnosed diabetes mellitus in 2009 in Iran was estimated as high as U.S.\$3.78 billion [12]. Therefore, prevention and early control of the risk factors associated with NCDs is a public health priority in Iran.

Childhood obesity is considered one of the major predisposing factors of most NCDs [13,14]. Therefore, having an insight on the prevalence and trend of childhood overweight and obesity can provide evidence-based information for health policymakers at national and international levels in order to implement programs for primordial and primary prevention of NCDs.

In recent years, various prevalence rates for childhood overweight and obesity have been reported; however, as Iran is the 18th largest country in the world, with an area of 1 648 195 km², and a population of around 75 million, it is necessary to condense the existing information to gain a general perception on this situation among Iranian children and adolescents. As the first report of its kind in MENA, the aim of this review was to assess the trend in the prevalence of childhood obesity and overweight in Iran, using systematic review and meta-analysis.

Material and methods

Search strategy

We searched multiple international (MEDLINE through PubMed and Scopus) and Iranian scientific databases (Scientific Information Database [SID] and IranMedex) for English and Persian language studies published between October 2007 and August 2012 that contained data on the prevalence of obesity and/or overweight among Iranian children and adolescents. The literature was searched combining keywords from the following concepts: the population (children, students, and adolescents and their related MeSH terms); the outcome (overweight, obesity, weight status, BMI, and anthropometric measures); the study design (prevalence, proportion, survey, descriptive, and epidemiology); and the study location (an English transcription of Iran and the name of Iranian medical universities) [15]. The search strategy used for PubMed and Scopus is shown in the supporting information (Table S1). For Iranian databases, Persian equivalents of the English keywords were applied, and the search strategy was constructed accordingly. Additionally, to maximize the sensitivity of the Persian literature search, we did not consider any time limitation. For studies published before October 2007, we entered all relevant articles from the previous meta-analysis on the prevalence of obesity in Iran [16]. Additionally, the reference list of all reviews and included papers was screened for any additional studies.

Study selection and eligibility criteria

Having removed duplicates, the relevant papers were selected in three phases. In the first and second phases, titles and abstracts of papers were screened, and irrelevant papers were excluded. In the last phase, the full text of identified papers was explored deeply to select only relevant papers. All three screening phases were done by two independent reviewers (RK and BS). Discrepancies were resolved by consultation and consensus.

In the next step, the eligibility of relevant papers was checked. We included studies if they were conducted among randomly selected healthy children/adolescents ages 2 to 18 y and provided data on the prevalence of obesity and/or overweight (including sample size) according to age–sex specific BMI cutoffs provided by the following organizations: U.S. Centers for Disease Control and Prevention in 2000 (CDC 2000) (i.e., 85th percentile BMI \leq 94th percentile was regarded as overweight and BMI \geq 95th percentile as obese) [17]; World Health Organization (WHO) (i.e., BMI \geq 2 z score was regarded as overweight and BMI \geq 3 z score as obese) [18]; or International Obesity Task Force (IOTF) (i.e., percentile curves that correspond to cutoff points of 25 kg/m² and 30 kg/m² for adult overweight and obesity, respectively) [19].

Papers were excluded if they reported a mixed prevalence of overweight and obesity, did not report the prevalence of obesity or overweight separately for children and adults, did not provide any prevalence data on obesity or overweight, or contained duplicate data. Duplicate data was identified by

examining similarities in the province in which the study was performed and by evaluating studies in terms of the author's names, the city name and rural/urban area, age and sex of participants, as well as the study year. Thus, when authors presented the results of the same study in more than one paper, the most recent one with the most comprehensive results and maximum sample size were included. However, for three studies, we extracted the most detailed information (on age–sex–specific prevalence) from five excluded articles. Moreover, we excluded the Persian version of a paper if it repeated results of the same English paper included in this review. We found one cohort study, Tehran Lipid and Glucose Study; for this cohort, we included one article, which presented the results of the two consecutive cross-sectional samples of this cohort [20].

The selection process of our systematic review is depicted in Figure 1.

Quality assessment

The methodological quality of each eligible paper was assessed using a checklist based on guidelines provided in an earlier study [21], in which papers were assigned up to a total score of 8 for fulfilling the main issues in prevalence studies, including sampling, measurements, and analysis.

Data extraction and abstraction

The required information that was extracted from all eligible papers was as follows: general characteristics of the study (first author's name, publication year, study year, study design, sampling method, and study location [i.e., province and city/town's name]), characteristics of the study population (age and sex of studied participants and the sample size), definition of obesity or overweight, and the prevalence of obesity or overweight by different definitions. Based on the information provided in each paper, we extracted the total prevalence and categorized it by age, sex, and age–sex combined. We analyzed the data in three age groups (2–6, 7–11, and 12–18 y).

One reviewer (RK) extracted the data, while another (BS) randomly selected 10% of the papers and checked their extracted data.

Statistical analysis

All eligible papers were entered into meta-analysis. The binomial distribution formula was used to compute the variance of point prevalence. Heterogeneity among studies was assessed using the Q statistic with a significance level of 0.1; I² was used to assess the degree of heterogeneity among studies. The random effects model was used to estimate the prevalence after heterogeneity was tested; if no heterogeneity was observed, the fixed model was used.

Separate meta-analyses were performed to estimate the prevalence of obesity and overweight overall, as well as age-, sex-, and age–sex specific. Moreover, to assess the prevalence trends, all meta-analyses were stratified by three categories of the study year. For the sake of space, forest plots were provided for the total and sex-specific prevalence.

We used meta-regression to assess the association of participant's age and sex and year of data collection with the prevalence of obesity and overweight. As the CDC criteria were the most frequent definition, our meta-regression was restricted to papers that used this definition. Statistical analyses were performed using Stata version 11.0, Texas, US.

Results

Study selection and characteristics

Our electronic search yielded 2 330 papers, of which 916 duplicate papers retrieved from more than one database were excluded. In the first (title evaluation) and second phase (abstract evaluation), 864 and 337 papers were excluded, respectively. Checking references and screening relevant studies from previous meta-analysis [16] resulted in 28 more papers being discovered. Finally, 241 papers were further assessed in full text; of these, 107 articles (49 English and 58 Persian) were included in this systematic review and meta-analysis. Figure 1 presents the flowchart of our study selection and the frequency of reasons for exclusion.

All studies were conducted among children or adolescents, except two, which were conducted among both children and adults, and two among participants younger than age 20 y. Five papers reported the prevalence in more than 1 y, and five were




Fig. 1. Flowchart of study selection. CDC, Centers for Disease Control and Prevention; IOTF, International Obesity Task Force; SID, Scientific Information Database; WHO, World Health Organization.

national surveys. Included studies were conducted between 1995 and 2010, the majority being conducted from 2005 to 2010. Thirteen studies used more than one definition to assess childhood overweight or obesity, with the CDC 2000 as the most frequently used definition. Age-, sex-, and age-sex-specific prevalence of overweight or obesity was found in 103, 97, and 90

papers, respectively. A total of 3 746 869 children and adolescents ages 2 to 18 y were included, with the interquartile range of sample sizes of 415 to 2076. The minimum and maximum quality scores of included studies were 3 and 8, respectively; about 57% had a score of more than 6 (Table S2 shows the characteristics of included papers [20,22–127]).

Table 1
Point estimate and 95% confidence interval for the prevalence of childhood obesity among Iranian children and adolescents using CDC definition criteria

	1995–1999		2000–2004		2005–2010	
	Number of studies (I ²)	Pooled estimate (95% CI)	Number of studies (I ²)	Pooled estimate (95% CI)	Number of studies (I ²)	Pooled estimate (95% CI)
All (y)	4 (99.3)	7.4 (0–14.9)	34 (95.4)	5.1 (4.4–5.8)	57 (97.5)	5.1 (4.9–5.4)
2–6	1	23.2 (21.1–25.2)	6 (84.5)	6.9 (4.8–9.1)	9 (97.0)	3.7 (3.4–3.9)
7–11	NA	NA	18 (95.8)	5.0 (4.0–5.9)	27 (97.7)	7.0 (5.5–8.5)
12–18	3 (82.3)	1.9 (0.2–3.5)	25 (96.0)	6.0 (4.9–7.1)	24 (95.8)	5.3 (4.2–6.4)
Girls (y)	4 (98.6)	6.9 (1.4–12.5)	29 (93.9)	4.8 (4.0–5.5)	40 (95.9)	5.7 (4.7–6.6)
2–6	1	21.9 (19.0–24.8)	5 (36.5)	5.3 (4.4–6.3)*	3 (96.5)	6.0 (0.1–11.9)
7–11	NA	NA	15 (93.7)	5.2 (4.0–6.3)	19 (96.4)	5.9 (4.2–7.6)
12–18	3 (82.3)	1.9 (0.3–3.5)	19 (95.9)	6.1 (4.7–7.6)	21 (92.5)	4.8 (3.8–5.8)
Boys (y)	1	24.4 (21.4–27.4)	22 (96.1)	5.8 (4.7–6.8)	28 (97.4)	7.5 (5.9–8.9)
2–6	1	24.4 (21.4–27.4)	5 (84.4)	5.4 (2.9–7.9)	3 (96.7)	8.6 (1.1–16.2)
7–11	NA	NA	12 (92.9)	4.7 (3.5–5.9)	14 (97.9)	8.2 (5.4–10.9)
12–18	NA	NA	14 (97.4)	6.8 (4.7–9.0)	14 (95.8)	5.9 (4.0–7.8)

CI, confidence interval; I², I square; NA, no published article was found which have examined the prevalence of childhood obesity in the selected years based on the determined definition

* Using fixed method.

Prevalence of childhood obesity and overweight

According to the CDC criteria, the overall prevalence of obesity remained constant during 2000 to 2004 and 2005 to 2010 and is estimated to be 5.1%. However, the sex-specific prevalence has increased during these periods of time. For girls, the prevalence of obesity increased from 4.8% between 2000 and 2004 to 5.7% between 2005 and 2010. The corresponding values for boys were 5.8% and 7.5%, respectively (Fig. S1). Comparison of the estimates of prevalence in different age groups showed that the overall prevalence of obesity has increased in children ages 2 to 6 and 7 to 11 y during the 2000s, but it has decreased among those ages 12 to 18 y. These findings were consistently observed in girls and boys (Table 1).

The overall prevalence of overweight has increased from 10.4% during 2000 to 2004 to 10.8% during 2005 to 2010, based on the CDC criteria, but the sex-specific prevalence showed a different pattern. In girls, the prevalence of overweight increased from 10.4% during 2000 to 2004 to 11.6% during 2005 to 2010. The corresponding values for boys were 10.2% and 9.4%, respectively (Fig. S2). Comparison of the prevalence estimates in different age groups showed that the overall prevalence of overweight has increased for children ages 7 to 11 y during the 2000s, but it has decreased among those ages 2 to 6 and 12 to 18 y. These findings were consistently observed for girls and boys (Table 2).

Compared with the CDC criteria, we found that the IOTF and WHO criteria tend to underestimate the prevalence of obesity and overestimate the prevalence of overweight (Tables S3 and S4). Additionally, due to the small number of studies performed before 2000, these results are only presented in figures (Figs. S1 and S2) and tables, and are not deeply discussed in this review.

Predicted prevalence of childhood obesity and overweight

The meta-regression analysis showed that the prevalence of overweight and obesity did not vary significantly with respect to year of data collection, sex, and age of study participants. Our findings showed that girls had a lower prevalence of obesity (difference, –1.2%, 95% confidence interval [CI], –2.7 to 0.3; $P = 0.123$) and higher prevalence of overweight (difference, 1.4%; 95% CI, –0.3 to 3.0; $P = 0.110$) compared with boys. Compared with participants ages 2 to 6 y, those ages 7 to 11 and 12 to 18 y had a lower prevalence of obesity (for 7–11 y, difference, –1.5%;

95% CI, –3.9 to 1.1; $P = 0.248$, and for 12–18 y, difference, –1.9%; 95% CI, –4.4 to 0.5; $P = 0.115$) and a higher prevalence of overweight (for 7–11 y, difference, 0.2%; 95% CI, –2.5 to 2.9; $P = 0.882$, and for 12–18 y, difference, 2.3%; 95% CI, –0.3 to 4.9; $P = 0.088$). Studies conducted from 1995 to 1999 revealed a higher prevalence of obesity (difference, 3.8%; 95% CI, –0.4 to 8.1; $P = 0.078$) and overweight (difference, 1.2%; 95% CI, –3.9 to 6.4; $P = 0.63$), and those during 2000 and 2005 showed a lower prevalence of obesity (difference, –0.4%; 95% CI, –1.9 to 1.1; $P = 0.588$) and overweight (difference, –0.1%; 95% CI, –1.7 to 1.6; $P = 0.911$), compared with studies that were conducted during 2005 and 2010. The trend in the predicted prevalence of obesity and overweight among different age–sex populations is presented in Table 3.

Discussion

This systematic review and meta-analysis has provided a precise estimate for the trend of obesity and overweight by age group and sex in Iranian children and adolescents. We documented various trends for overweight and obesity among different age groups of the Iranian pediatric population; in general, our findings revealed a higher increase in the trend of excess weight in children ages 2 to 6 and 7 to 11 y than in older age groups. This finding may suggest that younger children are getting fatter more rapidly than their older counterparts. This finding may reflect an epidemiologic transition along with a notable nutrition transition, in the Iranian population. The recent changes in lifestyle of children from a traditional style toward a Westernized pattern that includes consuming calorie-dense foods and engaging in sedentary activities might explain the higher prevalence of excess weight in younger than in older children. This finding is consistent with the situation reported in some other developing countries [6,128] and deserves attention at individual and public health levels.

Globally, a rapid rise has been reported for the prevalence of obesity and overweight among children ages < 5 y (i.e., 60% increase from 1990 to 2010, with an estimated rise of 9% from 2010 to 2020). It is noteworthy that the highest increase has been observed in low- and middle-income countries, with the Middle East being one of the regions with the highest prevalence of excess weight in preschoolers [129]. It should be noted that although part of the escalating trend of childhood obesity in poorer than in wealthier populations might be due to globalization and epidemiologic transition; however it could also be

Table 2

Point estimate and 95% confidence interval for the prevalence of childhood overweight among Iranian children and adolescents using CDC definition criteria

	1995–1999		2000–2004		2005–2010	
	Number of studies (I ²)	Pooled estimate (95% CI)	Number of studies (I ²)	Pooled estimate (95% CI)	Number of studies (I ²)	Pooled estimate (95% CI)
All (y)	3 (95.0)	10.9 (4.6–17.1)	33 (98.1)	10.4 (8.9–11.8)	50 (99.1)	10.8 (10.2–11.4)
<7	1	15.5 (13.7–17.3)	6 (95.0)	9.4 (5.1–13.7)	8 (99.8)	9.0 (7.9–10.1)
7–11	NA	NA	16 (98.1)	8.3 (6.3–10.3)	21 (97.3)	11.4 (9.4–13.3)
12–18	2	8.4 (3.4–13.3)	24 (94.8)	12.7 (11.1–14.3)	24 (93.4)	11.7 (10.3–13.2)
Girls (y)	3 (95.2)	11.5 (4.3–18.8)	28 (97.8)	10.4 (8.7–12.1)	39 (93.2)	11.6 (10.4–12.7)
<7	1	17.6 (14.9–20.2)	5 (91.1)	8.3 (4.3–12.3)	3 (84.4)	7.2 (3.8–10.5)
7–11	NA	NA	15 (97.7)	8.8 (6.5–11.1)	18 (95.2)	10.9 (8.8–12.9)
12–18	2	8.4 (3.4–13.3)	19 (97.5)	13.3 (10.4–16.1)	21 (89.6)	10.3 (10.7–13.8)
Boys (y)	1	13.3 (10.9–15.7)	21 (97.6)	10.2 (8.3–12.1)	27 (92.5)	9.4 (8.2–10.6)
<7	1	13.3 (10.9–15.7)	5 (97.2)	10.6 (3.4–17.8)	3 (59.2)	7.3 (5.1–9.5)
7–11	NA	NA	12 (94.8)	8.3 (6.4–10.1)	13 (96.1)	9.3 (6.9–11.7)
12–18	NA	NA	14 (97.2)	10.8 (7.9–13.6)	14 (92.2)	9.7 (7.7–11.8)

CI, confidence interval; I², I square; NA, no published article was found which have examined the prevalence of childhood obesity in the selected years based on the determined definition

due to the problem of stunting in children of developing countries. Therefore, the double burden of nutritional disorders and the persisting problem of micronutrient deficiency should be considered reasons for elevated BMI in children of developing countries.

Our meta-regression analysis showed that the prevalence of obesity and overweight did not vary significantly in girls and boys. We found that girls had a lower prevalence for obesity and a higher prevalence for overweight compared with boys. This may be because girls pay higher attention to their body image than boys [130]. Our findings show that, in general, the prevalence for obesity and overweight in Iranian children is not very high and the trend in its increase is not very sharp, but given that until a few years ago underweight was the main nutritional problem of Iranian children, the obtained trend in childhood obesity and overweight is alarming. We documented differences in the trend of obesity and overweight among Iranian children

and adolescents, but any kind of excess weight may have harmful health consequences. A remarkable point is that Asians have an ethnic predisposition to adverse health effects of obesity and overweight at lower BMI levels than Western populations [131]. Therefore, an increase in the BMI of children, even less than normal limits, may have long-term health consequences.

Another point that should be considered for children of developing countries is the problem of low birth weight, which by itself [132] or accompanied with a rapid growth spurt during childhood [133] makes one prone to NCDs. As low birth weight is not uncommon in such populations, including Iran [134], the escalating trends in excess weight of Iranian children may be an accumulating risk for those who were under weight at birth. Therefore, the true public health burden of childhood obesity in Iran may be higher than expected by the trend reported in our study.

Different definitions of childhood obesity make it hard to compare data between studies. We found that compared with the CDC criteria, the IOTF and WHO definitions tend to underestimate the prevalence of obesity and overestimate the prevalence of overweight among Iranian children. This finding is consistent with our previous nationwide study in showing close agreement of BMI cutoff points for Iranian children when using CDC criteria rather than other criteria [66].

Conclusion

This review, which is likely the first of its kind in the MENA region, appraised and synthesized a large number of studies. By calculating a pooled average result across studies, this study provides research evidence on the trend of obesity and overweight among Iranian children. The escalating trend of excess weight among young children is alarming and provides information for policymakers and health care providers at national and regional levels for preventive programs and interventions.

Acknowledgments

This study was funded by Isfahan University of Medical Sciences.

Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.nut.2013.08.011>.

Table 3

Trend in the predicted prevalence of obesity and overweight among Iranian children and adolescents 2 to 18 y based on CDC definition criteria

Gender	Age group (y)	Predicted prevalence (%)		
		1995–1999	2000–2004	2005–2010
Childhood obesity				
All	All	5.13	6.14	7.25
Girls	All	4.52	5.73	9.69
Boys	All	6.30	7.51	11.47
All	2–6	6.82	6.82	8.27
	7–11	6.33	6.33	7.78
	12–18	5.55	5.55	7.00
Boys	2–6	12.32	8.07	8.49
	7–11	10.85	6.6	7.02
	12–18	10.37	6.12	6.54
Girls	2–6	11.13	6.88	7.30
	7–11	9.66	5.41	5.83
	12–18	9.18	4.93	5.35
Childhood overweight				
All	All	10.43	11.25	10.88
Girls	All	10.81	11.29	12.18
Boys	All	9.63	10.11	11.10
All	2–6	9.07	9.67	9.07
	7–11	9.71	10.31	9.71
	12–18	11.83	12.43	11.83
Boys	2–6	9.83	8.50	8.59
	7–11	10.04	8.71	8.80
	12–18	12.13	10.80	10.89
Girls	2–6	11.18	9.85	9.94
	7–11	11.38	10.06	10.15
	12–18	13.48	12.15	12.24

References

- [1] Boutayeb A, Boutayeb S. The burden of non-communicable diseases in developing countries. *Int J Equity in Health* 2005;4:2.
- [2] Department of Public Information. Non-communicable diseases deemed development challenge of “epidemic proportions” in political declaration adopted during landmark general assembly summit, 2011. Available at <http://www.un.org/News/Press/docs/2011/ga11138.doc.htm>. Accessed March 10, 2013.
- [3] Finucane MM, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011;377:557–67.
- [4] Stevens GA, Singh GM, Lu Y, Danaei G, Lin JK, Finucane MM, et al. National, regional, and global trends in adult overweight and obesity prevalences. *Popul Health Metr* 2012;10:22.
- [5] Gupta N, Shah P, Nayyar S, Misra A. Childhood obesity and the metabolic syndrome in developing countries. *Indian J Pediatr* 2013;80(Suppl 1):S28–37.
- [6] Kelishadi R. Childhood overweight, obesity, and the metabolic syndrome in developing countries. *Epidemiol Rev* 2007;29:62–76.
- [7] Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047–53.
- [8] Mehio Sibai A, Nasreddine L, Mokdad AH, Adra N, Tabet M, Hwalla N. Nutrition transition and cardiovascular disease risk factors in Middle East and North Africa countries: reviewing the evidence. *Ann Nutr Metab* 2010;57:193–203.
- [9] Delavari A, Forouzanfar MH, Alikhani S, Sharifian A, Kelishadi R. First nationwide study of the prevalence of the metabolic syndrome and optimal cutoff points of waist circumference in the Middle East: the national survey of risk factors for noncommunicable diseases of Iran. *Diabetes Care* 2009;32:1092–7.
- [10] Haghdoost AA, Rezazadeh-Kermani M, Sadghirad B, Baradaran HR. Prevalence of type 2 diabetes in the Islamic Republic of Iran: systematic review and meta-analysis. *East Mediterr Health J* 2009;15:591–9.
- [11] Kelishadi R, Ardalan G, Gheiratmand R, Adeli K, Delavari A, Majdzadeh R. Paediatric metabolic syndrome and associated anthropometric indices: the CASPIAN Study. *Acta Paediatr* 2006;95:1625–34.
- [12] Javanbakht M, Baradaran HR, Mashayekhi A, Haghdoost AA, Khamseh ME, Kharazmi E, et al. Cost-of-illness analysis of type 2 diabetes mellitus in Iran. *PLoS One* 2011;6: e26864.
- [13] Li S, Chen W, Srinivasan SR, Xu J, Berenson GS. Relation of childhood obesity/cardiometabolic phenotypes to adult cardiometabolic profile: the Bogalusa Heart Study. *Am J Epidemiol* 2012;176(Suppl 7):S142–9.
- [14] Park MH, Falconer C, Viner RM, Kinra S. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obes Rev* 2012;13:985–1000.
- [15] Farhoudian A, Amini H, Sharifi V, Basirnia A, Mesgarpour B, Mansouri N. Systematic review of psychiatric disorders among Iranian adult general population. *Iran J Psychiatry* 2007;2:137–49.
- [16] Mirzazadeh A, Sadeghirad B, Haghdoost AA, Bahreini F, Rezazadeh Kermani M. The prevalence of obesity in Iran in recent decade: a systematic review and meta-analysis study. *Iran J Public Health* 2009;38:1–11.
- [17] Kuczumarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, et al. CDC growth charts: United States. *The Advance Data* 20001–20027.
- [18] World Health Organization. WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Suppl* 2006;450:76–85.
- [19] Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240–3.
- [20] Esmailzade A, Mirmiran P, Azadbakht L, Azizi F. Changes in prevalence of overweight among children and adolescents in Tehran: comparing three different definitions. *RJMS* 2006;13:19–30.
- [21] Boyle MH. Guidelines for evaluating prevalence studies. *Evid Based Ment Health* 1998;1:37–9.
- [22] Aazami M, Akbari M, Heshmati B, Alikhani M. The growth pattern among elementary school students in Kermanshah, Iran. *J Isfahan Med* 2012;30:1–8.
- [23] Ahmadi E, Tehrani AR, Ahmadi A. Prevalence of obesity, overweight and underweight among elementary school children in Southern Iran, 2009. *Am J Applied Sci* 2010;7:1439–42.
- [24] Ahmadi S, Shahsavari S, Ahmadi H, Tabatabaieifar T. Prevalence of overweight, obesity and underweight among high school students in Sanandaj: 2006–2007. *Iranian J Endocrinol Metab* 2010;12:153–9.
- [25] Akhavan Karbasi S, Fallah R, Golestan M, Sadr Bafghi M. Prevalence and risk factors of obesity and overweight among primary school children in Yazd. *J Shaheed Sadoughi Uni Med Sci & Health Serv* 2008;16:8–13.
- [26] Amanolahi A, Sohrabi MR, Montazeri A, Abazari AR, Kolahi AA. Study of obesity levels among school students. *Payesh J* 2012;11:83–9.
- [27] Amidi Mazaheri M, Hoseini M. Prevalence of underweight, overweight and obesity among high school girls in Isfahan. *Health Syst Res* 2010;6:1–6.
- [28] Amini M, Omidvar N, Kimiagar M. Prevalence of overweight and obesity among junior high school students in a district of Tehran. *J Res Med Sci* 2007;12:315–9.
- [29] Amirzadeh F, Agha-Alinejad H, Rajabi H, Sedjgh Sarvestani R. The relationship between overweight and obesity and economical status in Tehran girls students. *Res Sport Sci* 2004;3:69–85.
- [30] Asadi Noghani F. Prevalence of obesity and overweight among school children in Bandar Abbas, 2008. *J Hormozgan Univ Med Sci* 2011;15:218–26.
- [31] Asar S, Asghari S. Prevalence of obesity and overweight among 7–14 years old students in the city of Ahwaz. *Med J Ahvaz Univ Med Sci* 2005;44:11–20.
- [32] Ataei N, Hosseini M, Iranmanesh M. The relationship of body mass index and blood pressure in Iranian children <7 years old. *J Trop Pediatr* 2009;55:313–7.
- [33] Ayatollahi SMT, Mostajabi F. Prevalence of obesity among schoolchildren in Iran. *Obes Rev* 2007;8:289–91.
- [34] Azarbaijani MA, Tojari F, Habibinejad M. The relation between obesity, physical activity and socioeconomic status among girl students living in northern Tehran. *Fez J Kashan Univ Med Sci* 2011;15:132–8.
- [35] Basirat R, Abargouei AS, Esmailzadeh A. The association between household food insecurity and childhood obesity among Iranian school-aged children in Farokhshahr. *Koomesh J Semnan Univ Med Sci* 2012;13:254–63.
- [36] Baygi F, Eshraghian MR, Dorosty AR. Prevalence of obesity among school children in Neishabour (2005). *J Ghazvin Univ Med Sci* 2008;12:73–6.
- [37] Bazhan M, Kalantari N, Houshiar-Rad A. Pattern of fat distribution (waist to hip ratio) and its relationship with BMI among high school girls in Lahijan. *Iranian J Endocrinol Metab* 2006;8:163–8.
- [38] Behzadnia S, Vahidshahi K, Hamzeh Hosseini S, Anvari S, Ehteshami S. Obesity and related factors in 7–12 year-old elementary school students during 2009–2010 in Sari, Iran. *Med Glas (Zenica)* 2012;9:86–90.
- [39] Broughani M, Hamedinia MR, Assad MR, Zarei M. The study of the body mass index and physical activity among 11–14 year-old adolescent boys in Khooshab. *Knowledge Health J* 2010;5:12–8.
- [40] Dahri M, Safarian M, Hajifaraji M, Houshiar Rad A, Abadi AR. The association of overweight and obesity with menarche age and nutritional status in girls aged 11–15 in Mashhad. *Med J Mashhad Univ Med Sci* 2011;53:245–53.
- [41] Delvarianzadeh M, Khosravi A, Bolbolhaghghi N, Ebrahimi H. Relationship between anthropometric parameters with menarche age of school girls (11–14 years old) in Shahroud. *Knowledge Health J* 2008;3:43–7.
- [42] Doost Mohammadian A, Keshavarz A, Dorosty AR, Mahmoodi M, Sadrzadeh H. Comparison of body mass index and the intake of energy and some nutrients with reference values in female secondary school students in Semnan. *Payesh J* 2005;4:237–45.
- [43] Dorosty AR, Hojat P. The prevalence of obesity and its association with the parents obesity in primary school girls in Tehran in 2002. *Tehran Univ Med J* 2005;62:942–7.
- [44] Dorosty AR, Houshyar Rad A, Mohammad Pour Ahranjani B, Siassi F. Determination of the most relevant body mass index standard reference to define obese Iranian school age children. *Iranian J Nutr Sci Food Tech* 2009;4:71–80.
- [45] Dorosty AR, Siassi F, Reilly JJ. Obesity in Iranian children. *Arch Dis Child* 2002;87:388–91.
- [46] Ebrahimzadehkor, Kalantari N, Abadi AR. The prevalence of obesity and its relative factors among less than 5 years aged children, Bandar Turkmen district, Iran. *J Kerman Univ Med Sci* 2012;19:384–91.
- [47] Faghih S, Taleban F, Abadi A, Ansari N. Trends of overweight and obesity among 14–18 years old urban adolescent girls among senior high schools, Ahwaz, Iran. *Iranian J Diabetes Lipid Disord* 2009;9:19–26.
- [48] Fakhrzadeh H, pour Ebrahim R, Bandarian F, Heshmat R, Jalipour F, Nouri M, et al. Screening cardiovascular risk factors in high risk families: a study based the partnership model among students. *Iranian J Diabetes Lipid Disord* 2004;4:73–83.
- [49] Fesharakinia A, Zarban A, Sharifzadeh GR. Relationship of body mass index with serum lipids in elementary school students. *Indian J Pediatr* 2009;76:729–31.
- [50] Gaeni A, Kashef M, Samadi A, Fallahi A. Prevalence of underweight, overweight and obesity in preschool children of Tehran, Iran. *J Res Med Sci* 2011;16:821–7.
- [51] Gharakhanlou R, Farzad B, Agha-Alinejad H, Steffen LM, Bayati M. Anthropometric measures as predictors of cardiovascular disease risk factors in the urban population of Iran. *Arq Bras Cardiol* 2012;98:126–35.
- [52] Gharghare Chi R, Shoari N. Age of puberty and its relationship with body mass index in Iranian girls living in Tabriz. *Med J Tabriz Univ Med Sci* 2011;33:63–8.
- [53] Golestan M, Akhavan Karbasi S, Fallah Tafti M, Sharaf Aldini M. Prevalence of obesity, overweight and underweight in guidance school students. *J Shaheed Sadoughi Univ Med Sci Health Serv* 2008;16:31–5.
- [54] Haeri Behbahani B, Dorosty AR, Eshraghian MR. Assessment of obesity in children: fat mass index versus body mass index. *Tehran Univ Med J* 2009;67:408–14.

- [55] Haji Faraji M, Esfarjani F, Roustaei R, Khosh Fetrat MR, Kamrani Z. Assessment of anthropometric status of adolescents in public guidance schools in east Tehran, 2006. *Iranian J Nutr Sci Food Tech* 2007;2:37–43.
- [56] Hajian-Tilaki K, Heidari B. Prevalences of overweight and obesity and their association with physical activity pattern among Iranian adolescents aged 12–17 years. *Public Health Nutr* 2012;1:1–7.
- [57] Hajian-Tilaki KO, Sajjadi P, Razavi A. Prevalence of overweight and obesity and associated risk factors in urban primary-school children in Babol, Islamic Republic of Iran. *East Mediterr Health J* 2011;17:109–14.
- [58] Hosseini Kakhk SAR, Safari M, Hamedani N. Health-related factors in physical fitness in 12–14 year-old male adolescents in Sabzevar. *Iran J Sabzevar Univ Med Sci* 2011;18:55–66.
- [59] Hosseini M, Ataei N, Aghamohammadi A, Yousefifard M, Slimi SH, Ataei F. The relation of body mass index and blood pressure in Iranian children and adolescents aged 7–18 years old. *Iran J Public Health* 2010;39:126–34.
- [60] Hossieni M, Fallah H, Keshavarz SA. Comparison of nutritional status and related factors among pupils aged 11–14 year olds in public and private schools of Damghan (2005). *Koomeh Semnan Univ Med Sci* 2007;8:53–60.
- [61] Jafari Rad S, Keshavarz A, Dorosty AR. The nutritional factors by body mass index in young women in high school in Sari, 2003. *Iranian J Nutr Sci Food Tech* 2007;2:51–6.
- [62] Jamshidi I, Mohammad K, Jazayeri A, Hosseini S, Nouri Jalani K, Keshavarz A, et al. Association between body mass index and socioeconomic status in female students in Kish, Iran. *Payesh J* 2012;11:181–6.
- [63] Janghorbani M, Parvin F. Prevalence of overweight and thinness in high-school girls in Kerman, Iran. *Int J Obes Relat Metab Disord* 1998;22:629–33.
- [64] Javadi Nejad S, Karami M, Hashemnia N. Association between body mass index and dental development in 7–15 year old children in the city of Isfahan-Iran in the year 2008. *J Mashhad Dental School* 2010;34:109–16.
- [65] Karajibani M, Montazerifar F, Mohammadi M, Dashipour A. The prevalence of obesity and wasting in primary school girls in the city of Zahedan. *Tabib-e-Shargh* 2006;6:289–96.
- [66] Kelishadi R, Ardalan G, Gheiratmand R, Majdzadeh R, Hosseini M, Gouya MM, et al. Thinness, overweight and obesity in a national sample of Iranian children and adolescents: CASPIAN Study. *Child Care Health Dev* 2008;34:44–54.
- [67] Kelishadi R, Hashemipour M, Sarrafzadegan N, Sadri GH, Ansari R, Alikhani H, et al. Obesity and associated modifiable environmental factors in Iranian adolescents: Isfahan Healthy Heart Program – Heart Health Promotion from Childhood. *Pediatr Int* 2003;45:435–42.
- [68] Khabaz Khoob M, Fotouhi A, Hashemi H, Mohammad K. Obesity and overweight in Tehran, 2002. *Iranian J Diabetes Lipid Disord* 2008;7:315–24.
- [69] Khabaz Khoob M, Fotouhi A, Mohammad K. Thinness and obesity based on body mass index in Dezfool schoolchildren in 2004. *Iranian J Epidemiol* 2007;3:35–43.
- [70] Khaji A, Khodaei S, Karbakhsh M, Faeghi A, Azizi S, Firoozian A, et al. Blood pressure and obesity in young adolescents in Tehran. *Iran J Pediatr* 2006;16:45–50.
- [71] Khosh Fetrat MR, Rahmani K, Kalantari N, Ghaffarpour M, Mehrabi Y, Esmaeilzade A. Assessment and comparison of food consumption patterns and body mass index of adolescent boys in urban and rural areas of Zarrinshahr, Isfahan. *Payesh J* 2007;6:119–27.
- [72] Maddah M, Nikooyeh B. Obesity among Iranian adolescent girls: location of residence and parental obesity. *J Health Popul Nutr* 2010;28:61–6.
- [73] Maddah M, Nikooyeh B. Factors associated with overweight in children in Rasht, Iran: gender, maternal education, skipping breakfast and parental obesity. *Public Health Nutr* 2010;13:196–200.
- [74] Maddah M, Rashidi A, Mohammadpour B, Vafa R, Karandish M. In-school snacking, breakfast consumption, and sleeping patterns of normal and overweight Iranian high school girls: a study in urban and rural areas in Guilan. *Iran J Nutr Educ Behav* 2009;41:27–31.
- [75] Mehrabi E, Kargarfard M, Kelishadi R, Mojtahedi H. Effects of obesity on pulmonary function in obese, overweight, and normal students. *J Isfahan Med School* 2012;30:1–9.
- [76] Mirhosseini NZ, Yusoff NA, Shahar S, Parizadeh SM, Mobarhen MG, Shakery MT. Prevalence of the metabolic syndrome and its influencing factors among adolescent girls in Mashhad, Iran. *Asia Pac J Clin Nutr* 2009;18:131–6.
- [77] Mirmohammadi S, Hafezi R, Mehrparvar AH, Rezaeian B, Akbari H. Prevalence of overweight and obesity among Iranian school children in different ethnicities. *Iran J Pediatr* 2011;21:514–20.
- [78] Mirzaei M, Karimi M. Prevalence of overweight and obesity among the first grade primary students in Yazd. *J Ilam Univ Med Sci* 2010;18:43–9.
- [79] Mirzaeian S, Fakhari M, Hosseini R, Hassanzadeh A, Esmaeilzadeh A. Association between duration of breastfeeding and subsequent overweight and obesity in female adolescents. *Iranian J Nutr Sci Food Tech* 2011;6:65–74.
- [80] Moadab MH, Kelishadi R, Hashemipour M, Amini M, Poursafa P. The prevalence of impaired fasting glucose and type 2 diabetes in a population-based sample of overweight/obese children in the Middle East. *Pediatr Diabetes* 2010;11:101–6.
- [81] Moayeri H, Bidad K, Aghamohammadi A, Rabbani A, Anari S, Nazemi L, et al. Overweight and obesity and their associated factors in adolescents in Tehran, Iran, 2004–2005. *Eur J Pediatr* 2006;165:489–93.
- [82] Moghadasi M, Naser K, Ghanbarzadeh M, Shakerian S, Razavi A. Prevalence of overweight, obesity and physical fitness in Shiraz adolescents. *Iranian J Endocrinol Metab* 2011;12:476–82.
- [83] Mohammad Pour Kolde M, Fouladvand MA, Avakh Keysami M. Prevalence of overweight and obesity among Bushehr high-school girls at aged 14–17 years old. *Iranian South Med J* 2012;15:221–32.
- [84] Mohammadian S, Khoddam H, Kaveh M. Related factors of obesity and overweight among secondary school girls (Gorgan-Iran). *J Gorgan Univ Med Sci* 2010;12:57–63.
- [85] Mohammadpour-Ahranjani B, Rashidi A, Karandish M, Eshraghian MR, Kalantari N. Prevalence of overweight and obesity in adolescent Tehrani students, 2000–2001: an epidemic health problem. *Public Health Nutr* 2004;7:645–8.
- [86] Mohkam M, Karimi A, Eslami N, Khatami A, Fallah F, Maham S, et al. Blood pressure screening in school-aged children in Tehran. *Iran J Kidney Dis* 2011;5:229–33.
- [87] Mojibian M, Ghilian Z. Prevalence of obesity among women in Yazd city in 2000. *J Shaheed Sadoughi Univ Med Sci Health Serv* 2001;9:36–42.
- [88] Montazerifar F, Karajibani M, Dashipour A. The prevalence of obesity and wasting and their correlation with food intake in female junior school students in Zahedan, 2004. *Iranian South Med J* 2006;9:66–76.
- [89] Montazerifar F, Karajibani M, Rakhshani F, Hashemi M. Prevalence of underweight, overweight and obesity among high-school girls in Sistan va Baluchistan. *East Mediterr Health J* 2009;15:1293–300.
- [90] Mortazavi Z, Roudbari M. Breakfast consumption and body mass index in primary, secondary and high school boys in Zahedan 2005–2006. *Iranian J Endocrinol Metab* 2010;12:345–51.
- [91] Mosavi Jazayeri SMH. Overweight and obesity among school-aged children of Metropolitan Tehran, Iran. *Pak J Nut* 2005;4:342–4.
- [92] Mostafavi H, Dabagh Manesh MH, Zare N. Prevalence of obesity and overweight in adolescents and adult population in Shiraz. *Iranian J Endocrinol Metab* 2005;7:57–66.
- [93] Motlagh ME, Kelishadi R, Ziaoddini H, Mirmoghtadaee P, Poursafa P, Ardalan G, et al. Secular trends in the national prevalence of overweight and obesity during 2007–2009 in 6-year-old Iranian children. *J Res Med Sci* 2011;16:979–84.
- [94] Mozaffari H, Nabaei B. Obesity and related risk factors. *Indian J Pediatr* 2007;74:265–7.
- [95] Mozaffari Khosravi H, Hossein Zadeh M, Shariati Bafghi SE, Mozaffari Khosravi V. Prevalence of eating disorders and obesity among highschool girls in Yazd, 2010–2011. *Tolooeh Behdasht J* 2011;10:38–49.
- [96] Nabavi M, Karimi B, Ghorbani R, Mazloom Jafarabadi M, Talebi M. Prevalence of obesity and some related factors among students aged 7 to 12 in Semnan, Iran. *Payesh J* 2010;9:443–51.
- [97] Nakhostin-Roohi B, Sojudi A, Niknam Z. Nutritional status and physical activity in 7–11 year-old female students of Pars-Abad in the province of Ardabil. *Medicina dello Sport* 2010;63:521–9.
- [98] Nemati A, Barak M, Naghizadeh Baghi A, Abbasgholizadeh N, Homapour F, Hazrati S, et al. Relationships between anthropometrical indices and socio-economic differences for children at 6 years old living in urban areas of Ardebil. *Iran J Applied Sci* 2008;8:3748–52.
- [99] Pour Aram H, Aminpour A, Kalantari N, Golestan B, Abtahi M. Survey of anthropometrics indexes weight, height, TSF and BMI status of adolescent girls in Astaneh Ashrafieh, Gilan 2000. *J Guilan Univ Med Sci* 2003;12:56–66.
- [100] Pour Ghasem Gargari B, Hamed-Behzad M, Ghassab Pour S, Ayat A. Prevalence of overweight and obesity among high-school girls in Tabriz, Iran, in 2001. *Food Nutr Bull* 2004;25:288–91.
- [101] Pour Ghasem Gargari B, Hamed-Behzad M, Seyed Sajadi N, Koushavar D, Karami S. Relation of body mass index to eating attitude in Tabrizian high school girls. *Med J Tabriz Univ Med Sci* 2010;32:24–9.
- [102] Rafraf M, Gargari BP, Safaiyan A. Prevalence of prehypertension and hypertension among adolescent high school girls in Tabriz, Iran. *Food Nutr Bull* 2010;31:461–5.
- [103] Rahmani Nia F, Daneshmandi H, Darbani H. Determination of overweight and obesity prevalence among male students and its relationship with physical activity-level. *Harkat J* 2005;23:47–60.
- [104] Rahmani Nia F, Daneshmandi H, Taghipour A. The relationship between underweight and overweight to physical fitness and socioeconomic status of students. *J Sport Biosci* 2009;1:127–44.
- [105] Rahmanpour H, Heidari R, Mousavinasab SN, Sharifi F, Fekri S. The prevalence of polycystic ovarian syndrome in 14–18 year old girls of Zanjan high school, 2008. *J Zanjan Univ Med Sci Health Serv* 2009;17:79–88.
- [106] Rouhani MH, Mirseifinezhad M, Omrani N, Esmaeilzadeh A, Azadbakht L. Fast food consumption, quality of diet, and obesity among Isfahanian adolescent girls. *J Obes*; 2012:597924.
- [107] Sadeghi M, Lynch CD, Arsalan A. Is there a correlation between dental caries and body mass index-for-age among adolescents in Iran? *Community Dent Health* 2011;28:174–7.

- [108] Saffari F, Jalilolghadr S, Esmailzadehha N, Shariatinejad K. Anthropometric indices in 6–14 year-old girls in Qazvin. Iran. *J Isfahan Med School* 2012;29:1–11.
- [109] Salem Z, Vazirinejad R. Prevalence of obesity and metabolic syndrome in adolescent girls in South East of Iran. 2007. *Pak J Med Sci* 2009;25:196–200.
- [110] Salem Z, Vazirinejad R. Prevalence of the metabolic syndrome components among 7–11 year-old children in Rafsanjan, 2008. *Iranian J Nutr Sci Food Tech* 2010;5:63–71.
- [111] Shahidi N, Mirmiran P, Amirkhani F. The prevalence of obesity and abdominal obesity and its association with nutritional habits among high school adolescent boys in Tabriz city. *Sci J Shahid-Beheshti Univ Med Sci* 2004;28:255–63.
- [112] Shajari H, Shajari A, Sepahi MA, Mehrparvar AH, Roghani R, Nakhaei MH. Relationship between arterial blood pressure and body mass index of school age children of southern region of Iran. *Acta Med Iran* 2011;49:737–41.
- [113] Soheili Far J, Emdadi M. Relation between breast feeding with overweight and obesity in Hamadan primary school children. *Sci J Hamadan Univ Med Sci Health Serv* 2005;12:54–7.
- [114] Sotoudeh G, Khosravi S, Khajehnasiri F, Khalkhali HR. High prevalence of overweight and obesity in women of Islamshahr, Iran. *Asia Pac J Clin Nutr* 2005;14:169–72.
- [115] Tabatabaei M, Dorosty AR, Siassi F, Rahimi A. Using different reference values to determine prevalence of obesity among school children in Ahvaz. *J School of Public Health Inst Public Health Res* 2004;2:11–8.
- [116] Taheri F. Epidemiological study of obesity in school age children in Birjand. *J Birjand Univ Med Sci* 2002;9:18–22.
- [117] Taheri F, Kazemi T. Prevalence of overweight and obesity in 7 to 18 year-old children in Birjand/Iran. *Iran J Pediatr* 2009;19:135–40.
- [118] Vafa MR, Afshari S, Moslehi N, Salehpoor A, Hosaini F, Goharinezhad M, et al. Relationship between infant nutrition feeding and childhood obesity in first grade Tehranian students of primary schools, 2009. *Iranian J Endocrinol Metab* 2011;12:505–12.
- [119] Vahidi Nia AA, Sardarian K. Assessment of nutritional status in rural regions of Hamadan city/Iran, 2003. *Iranian South Med J* 2005;8:187–93.
- [120] Valizadeh S, Jabraili M, Ghojzadeh M, Kasraie B. The relationship between leisure activities pattern and body mass index in girl students in elementary schools in Tabriz. *Nurs Midwifery J* 2010;5:54–63.
- [121] Veghari G. Prevalence of overweight, obesity and socio-demographic related factors among Iranian northern school children. *J Bio Sci* 2011; 11:487–91.
- [122] Veghari G, Mohsen S. The comparative study of body mass index distribution among preschool children in a 7 years periods in north of Iran. *J Applied Sci* 2007;7:2681–5.
- [123] Yusefi Z, Rajaei N. Assessment of the relationship of anthropometric parameters and intelligence quotient to age of menarche (Mashhad, Autumn 2000). *Iranian J Obstet Gyn Infertil* 2000;56:21–9.
- [124] Zakeri M, Sedaghat M, Motlagh ME, Tayari Ashtiani R, Ardalan G. BMI correlation with psychiatric problems among 10–18 years Iranian students. *Acta Med Iran* 2012;50:177–84.
- [125] Zamani M, Farimani M, Zareian Z. The prevalence of oligomenorrhea and its relation to obesity in high school girls in Jahrom. *Iranian J Obstet Gyn Infertil* 2012;15:36–41.
- [126] Zareie M, Hamedinia MR, Haghghi AH, Chamari M, Broughani M. The epidemiology of obesity and underweight and their associations with physical activity and diet patterns among 12–14 years-old adolescent boys in Sabzevar, Iran. *Payesh J* 2011;10:243–53.
- [127] Ziaei Kajbaf T, Asar S, Alipoor MR. Relationship between obesity and asthma symptoms among children in Ahvaz, Iran: a cross sectional study. *Ital J Pediatr* 2011;37:1.
- [128] Núñez-Rivas HP, Monge-Rojas R, León H, Roselló M. Prevalence of overweight and obesity among Costa Rican elementary school children. *Rev Panam Salud Publica* 2003;13:24–32.
- [129] de Onis M, Blossner M, Borghi E. Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr* 2010;92: 1257–64.
- [130] Kelishadi R, Marashinia F, Heshmat R, Motlagh ME, Qorbani M, Taslimi M, et al. First report on body image and weight control in a nationally representative sample of a pediatric population in the Middle East and North Africa: the CASPIAN-III study. *Arch Med Sci* 2013;9:210–7.
- [131] Nubé M. The Asian enigma: predisposition for low adult BMI among people of South Asian descent. *Public Health Nutr* 2009;12:507–16.
- [132] Newsome CA, Shiel AW, Fall CH, Phillips DI, Shier R, Law CM. Is birth weight related to later glucose and insulin metabolism? A systematic review. *Diabetic Med* 2003;20:339–48.
- [133] Nobili V, Alisi A, Panera N, Agostoni C. Low birth weight and catch-up-growth associated with metabolic syndrome: a ten year systematic review. *Pediatr Endocrinol Rev* 2008;6:241–7.
- [134] Jafari F, Eftekhar H, Pourreza A, Mousavi J. Socio-economic and medical determinants of low birth weight in Iran: 20 years after establishment of a primary healthcare network. *Public Health Nutr* 2010;124:153–8.